



Impacts of biological nitrogen fixation on N₂O emissions from global natural terrestrial ecosystem soils: An Analysis with a process-based biogeochemistry model

5 Tong Yu¹ and Qianlai Zhuang^{1,2}

¹Earth, Atmospheric, and Planetary Sciences, Purdue University, West Lafayette IN 47907, USA

²Department of Agronomy, Purdue University, West Lafayette, IN 47907, USA

*Correspondence to: *Qianlai Zhuang (qzhuang@purdue.edu)*

10

15

20



25 **Abstract.** Biological nitrogen fixation plays an important role in the global nitrogen cycle. However, the fixation
rate has been usually measured or estimated at a particular observational site. To quantify the fixation amount at the
global scale, a process-based model is needed. This study develops a biological nitrogen fixation model and couples
it with an extant biogeochemistry model of N₂O emissions to examine the fixation rate and its effects on N₂O
emissions. The revised N₂O emission model better matches the observed data in comparison with our previous
30 model that has not considered the fixation effects. The new model estimates that tropical forests have the highest
fixation rate among all ecosystem types, and decrease from the equator to the polar region. The estimated nitrogen
fixation in global terrestrial ecosystems is 61.5 Tg N yr⁻¹ with a range of 19.8 - 107.9 Tg N yr⁻¹ in the 1990s. Our
estimates are relatively low compared to some early estimates using empirical approaches, but comparable to more
recent estimates that involve more detailed processes in their modeling. Furthermore, we estimate that the fixation
35 contributes to -5% to 20% changes in N₂O emissions compared to our previous estimates, depending on ecosystem
types and climatic conditions. This study highlights that there are relatively large effects of the biological nitrogen
fixation on ecosystem nitrogen cycling and soil N₂O emissions and calls for more comprehensive understanding of
biological nitrogen fixation and more observational data for different ecosystems to improve future quantification of
the fixation and its impacts.

40 1. Introduction

In most terrestrial ecosystems, nitrogen (N) available for plants is generally limited although it is the most
abundant element in the atmosphere (LeBauer and Tresder, 2008). It usually enters terrestrial ecosystems through
processes of N deposition and from biological N fixation (BNF). Nitrogen deposition is a physical process,
representing the direct input of reactive nitrogen (organic N, ammonia, and nitrogen oxides (NO_x) including nitric
oxide (NO), nitrogen dioxide (NO₂), nitric acid (HNO₃) and organic nitrates) from the atmosphere to biosphere.
45 BNF, a biochemical process that converts nonreactive nitrogen (N₂) to reactive nitrogen, provides a liaison between
the atmosphere and biological systems. Lightning is also a way to convert N₂, adding 3-5 Tg N yr⁻¹ to terrestrial
ecosystems (Levy and Moxim, 1996). Nitrogen input via rock weathering is another important source for terrestrial
ecosystems, adding 3-10 kg N ha⁻¹ yr⁻¹ (Morford et al., 2011; Houlton et al., 2018). BNF is significantly greater than
50 lightning induced N fixation (Galloway et al., 1995). Taken together, natural N fixation is the primary source in the
absence of human activities to global terrestrial ecosystems. For natural terrestrial ecosystems, the amount of N
added is approximately balanced by the nitrogen converted back to the atmosphere (Stedman and Shetter, 1983) and
lost into ocean and other aquatic systems.

Once entering terrestrial ecosystems, N can be taken up by plants and microbes, and converted into other
55 oxidized forms through mineralization, nitrification and denitrification. In terrestrial ecosystems, N fixation
generally affects the nitrogen cycle and nutrient level to constrain plant productivity. Any change of nitrogen input
to terrestrial ecosystems will influence their soil N₂O emissions. Atmospheric N₂O absorbs infrared wave and reacts
with ozone, affecting the global climate and atmospheric chemical conditions. On a per-unit-weight basis, N is
almost 300 times more effective in trapping heat than carbon dioxide. Its atmospheric concentrations have increased
60 from 286 ppb in 1950 to 327 ppb in 2015 observed in Cape Grim, Australia by the Advanced Global Atmospheric



Gases Experiment (AGAGE) Network, and in Alaska, Hawaii and South Pole by National Oceanic and Atmospheric Administration (NOAA). While the increase of concentration is significant due to rising industrial and agricultural emissions, natural ecosystems contribute about 60% of the increase (EPA, 2010). To date, the quantification of the trend and magnitude of N₂O emissions from natural terrestrial ecosystems is still with large uncertainties.

65 Furthermore, the effect of BNF in natural ecosystems on N₂O emissions has not been evaluated.

In the process of BNF, N₂ is converted to ammonia by certain soil microorganisms which can then be utilized by and incorporated into plants. In natural environment, N fixation is conducted by two types of microorganisms: asymbiotic organisms, including blue-green algae, lichens and free-living soil bacteria (Belnap, 2002; Granhall & Lid-Torsvik, 1975), and symbiotic organisms, such as fungi and nodule forming Rhizobium species. Among them,
70 the most dominant fixers are leguminous plants and their N fixation mechanisms are also best known (Sullivan et al., 2014; Vitousek et al., 2013). To date, the amount of N fixation by legumes is estimated in the range of 11.3-33.9 kg N ha⁻¹ yr⁻¹ (2.8-8.4 g m⁻² yr⁻¹) in natural terrestrial ecosystems. A symbiotic relationship exists between legume plants and bacteria. In the process, legume plants provide the bacteria energy through photosynthesis, meanwhile, the bacteria around the rhizobia-supply the legume N in the form of ammonia.

75 This study first models the BNF from the symbiotic relationship between legume plants and bacteria. The model is then coupled with an existing N₂O biogeochemistry to quantify the BNF impact on N₂O emissions from global terrestrial ecosystem soils.

2. Methods

2.1 Overview

80 We first develop a BNF model and then couple the model with an earlier version of N₂O biogeochemistry model (Yu and Zhuang, 2019). The revised model is then used to quantify the BNF at the global scale and its impacts on soil N₂O emissions from natural terrestrial ecosystems. The BNF rate estimates consider the effects of environmental conditions including temperature, soil moisture, soil mineral nitrogen content and soil carbon content. The new model is calibrated and evaluated with observed N fixation rate data from published studies for various
85 natural terrestrial ecosystems from the Arctic to tropical ecosystems. The model sensitivity to model input is analyzed. The model is then extrapolated to the global terrestrial ecosystems at a monthly step and a spatial resolution of 0.5° by 0.5° for the final decade of the 20th century. The effects of BNF and atmospheric nitrogen depositions on N₂O emissions from natural terrestrial ecosystem soils are analyzed.

2.2 Model Description

90 The Terrestrial Ecosystem Model (TEM) is a process-based model that simulates carbon and nitrogen dynamics, hydrological and thermal processes for terrestrial ecosystems. Although many efforts were made to incorporate more details of the N cycle, the N input from the atmosphere to ecosystems has not fully been incorporated to date, especially the BNF as input. Here we improve the N dynamics within TEM by considering N



95 fixation by legumes. The other calculations of carbon and N fluxes and pools are inherited from an earlier version of TEM (Zhuang et al., 2003; Yu and Zhuang, 2019).

BNF is the most significant process in either symbiotic or non-symbiotic forms converting stable molecular N₂ into N chemical compounds that are available to plants. For most terrestrial ecosystems, N fixer could be in many forms, such as free-living bacteria, lichens, and blue algae. But among them, symbiotic BNF is a dominant process to provide biologically accessible N, and most systematical BNF is regulated by legume plants, especially in
100 cropland and semi-natural environment (Mus et al., 2016). In natural environment, contribution from legume can be significant but with large uncertainties, which is greatly determined by various environmental conditions (Lindemann and Glover, 1996). In this study, the N fixation via legume plants is modeled considering (1) the accessible N concentration in soils, (2) the limitation of temperature, (3) soil water status, (4) the carbon demand for N fixation, and for spatial simulation, (5) the percentage of N fixing plants for each ecosystem type as:

$$105 \quad N_{fix} = N_{fixpot} f_t f_w f_N f_c f_{plant} \quad (1)$$

where N_{fixpot} is the potential N fixation rate (g N day⁻¹), f_t is the influence function of soil temperature, f_w is the soil water function, f_N is the function of root substrate N concentration, f_c is the function of plant carbon availability, and f_{plant} is the function of legume plant coverage.

The potential N fixation is highly related to the total N demand of plants and the available nitrogen in soils.
110 Theoretically, the definition of potential N fixation rate should be the difference between the demand and supply of N. Both of them vary with plant types, stages of growth and soil conditions. For large spatial-scale simulations for various ecosystem types, it is impossible to derive potential N fixation because of data availability. N_{fixpot} can be estimated based on dry matter of root, nodule or plant dry matter (Voisin et al, 2003, 2007). However, root biomass is also difficult to measure directly. In most published studies, the potential nitrogen fixation rate was measured
115 using an acetylene reduction array (ARA) method (Hardy et al, 1968, 1973), and some used ¹⁵N methods (Shearer and Kohl, 1986). In our simulation, N_{fixpot} is assumed to be a constant for each ecosystem type. The N_{fixpot} range is determined from literature and specific values for various ecosystem types are obtained through model parameterization.

Soil temperature is a controlling factor for both microbial activities and plant growth. A large number of
120 studies show that different plants have slightly different preferences for temperature (Montanez et al, 1995; Breitbarth et al., 2007; Gundale et al., 2012). For soybean, 20-35 °C is optimal (Boote et al., 2008), and for white clover the optimal temperature can be 13-26 °C (Wu and McGechan, 1999). The activity of microbes responds slightly differently to temperature among species. For most of them, the optimum temperature is 20-25 °C, and at 12-35 °C the activity is not limited. Generally, BNF increases as the temperature rises from minimum temperature
125 (0-5 °C) for N fixation to optimal temperature, maximum rate occurs within an optimal range (15-25°C), and decreases from optimal to maximum temperature above which BNF will stop at 35-40 °C:



$$f_t = \begin{cases} 0 & \text{when } (t < t_{\min} \text{ or } t > t_{\max}) \\ \frac{t-t_{\min}}{t_{\text{optL}}-t_{\min}} & \text{when } (t_{\min} \leq t < t_{\text{optL}}) \\ 1 & \text{when } (t_{\text{optL}} \leq t \leq t_{\text{optH}}) \\ \frac{t_{\max}-t}{t_{\max}-t_{\text{optH}}} & \text{when } (t_{\text{optH}} < t \leq t_{\max}) \end{cases} \quad (2)$$

130 where the upper limit (t_{\max}) is set to 45 °C. There is no lower limit, but when t is low enough, f_t will be close to zero (Wu and McGechan, 1999; Boote et al., 2008; Holzworth et al., 2014) (Table 1). For the convenience in computing, a lower limit is set in our model. When the temperature goes beyond its upper or lower limit, f_t is assumed to be 0.

Water stress has a direct effect on nitrogen fixing system (Sprent, 1972). With proper temperature, soil moisture condition is the major factor controlling nitrogen fixation rate (Srivastava and Ambasht, 1994). Soil water deficit and flood dramatically inhibits N fixation because of drought stress and oxygen deficit, respectively (Omari et al., 2004; Mario et al., 2007). In our model, the water factor is linearly related with soil water content (Williams, 1990; Wu and McGachan, 1999):

$$f_w = \begin{cases} 0 & \text{when } (W_f \leq W_a) \\ \varphi_1 + \varphi_2 & \text{when } (W_a < W_f < W_b) \\ 1 & \text{when } (W_f \geq W_b) \end{cases} \quad (3)$$

140 where W_f (J kg^{-1}) is the available soil water, which is defined as the ratio of water content to that at the field capacity. In soils, water potential generally includes osmotic and matrix potentials, ranging from -0.1 to -0.3 bar for typical soils, which has little effects on the N fixation. But when the soil gets very dry, the potential can be up to -100 to -200 bar and increases rapidly. W_a is the bottom threshold below which N fixation is totally restricted by soil moisture. W_b is the upper threshold above which nitrogen fixation is not limited by soil moisture. φ_1 and φ_2 are parameters representing the linear relationship between soil water content and its effect on N fixation (Table 1).

145 It is generally thought that more substrate N in soils will slow down the N fixation, because plants can uptake N directly from soil with less energy (Vitousek and Field, 1999). By comparison, N fixation needs more energy and consumes more carbon than plant N uptake does. Thus, the N fixation is only considered to occur when the direct N uptake from soil cannot meet the plant N demand. In our model, the inhibition effect of N is defined as (Wu and McGehan, 1999):

$$150 \quad f_N = \begin{cases} 1 - f_{Nup} \ln(1000 - N_s) & \text{when } (N_s \geq 0.001) \\ 1 & \text{when } (N_s < 0.001) \end{cases} \quad (4)$$

Where f_{Nup} is a parameter related to legume biological N fixation and soil N. N_s is the soil mineral N (g N m^{-2}). BNF efficiency shows a natural logarithmic relation with the soil mineral N.

155 N fixers get photosynthetic carbohydrate support from plants. Because the product of every unit of nitrogen consumes a certain amount of carbon, the lack of carbon supply will inhibit the N fixation. The carbon cost for per



160

unit of fixed N varies widely depending on environmental conditions and ecosystem types. For example, the consumption of carbon is only 1.54 times of fixed N for cowpea (Layzell et al., 1979), and it can be 6.3 to 6.8 times for soybeans (Ryle et al., 1979). It is also related to the life cycle of plants. The carbon effect is modeled following a Michaelis-Menten equation (Boote et al., 1998):

$$f_c = \frac{1}{1 + K_c / C_r} \quad (5)$$

where C_r is the soil carbon content (g C m^{-2}) to represent carbon availability from plants to N fixers. K_c is the Michaelis-Menten constant, which is plant species dependent.

2.3 Data

165

The classification of land cover and leguminous biomes were derived from the combination of the International Geosphere and Biosphere (IGP) land-cover classification system and the study of Schrire et al (2005). The experimental N_2 fixation data for model calibration were collected for 7 major ecosystem types (Figure 1). Nitrogen fixation rates were determined with acetylene reduction assay (ARA) method in most published studies (Table 2, data were from Cleveland et al. (1999)), expressed in $\text{kg N m}^{-2} \text{yr}^{-1}$. Some of them were measured with the ^{15}N natural abundance technique. Measurements of N_2O emissions affected by legumes were collected at 8 sites, including tropical rainforest in Brazil and Peru, tropical dry forests in Puerto Rico, temperate forest in New York, grassland in Germany and Scotland, and pastures in Australia and Costa Rica (Table 3). Monthly N_2O emission data were organized for comparison with model simulations.

170

175

The parameters for N_2 fixation module were initialized with *a priori* values (Table 2). Ecosystem-specific and microbe guild-specific parameters were inherited from previous TEM model (Zhuang et al., 2003; Yu and Zhuang, 2019). The global simulations were conducted at a spatial resolution of 0.5 by 0.5 degree and at a monthly time step. Historical climate data including temperature, precipitation, cloudiness and water vapor pressure were derived from the Climate Research Unit (CRU) (Mitchell and Jones, 2005). Soil texture data were from Melillo et al. (1993) and Zhuang et al. (2003). Other initial conditions including vegetation properties, soil carbon content and soil nitrogen contents were from Chen and Zhuang (2013) and Zhuang et al. (2012).

180

For regional simulations, the total amount of fixed N is also influenced by legume coverage. For each ecosystem type, we estimated the coverage according to the distribution of legume plants and field studies (Table 4, the coverage data are compiled from Cleveland et al., (1999)), where the minimum and maximum values were derived from the abundance of N-fixers.

185

2.4 Model calibration and site-level validation

Most model parameters are legume-specific or vegetation-specific and are adjusted based on value ranges from previous studies (Table 1). Model is parameterized for 11 representative natural terrestrial ecosystems (Table 2). Root mean square error (RMSE) and coefficient of determination ($0 \leq R^2 \leq 1$) were used for model calibration. RMSE



190 was calculated to show the mean difference between simulated data and observational values. The model is iterated
with changing parameters until the RMSE reached a certain value for each site. Most parameters in the model
driving nitrification and denitrification have been defined and calibrated in previous studies (Yu and Zhuang, 2019).
The calibrated model is evaluated at the site level (Table 3) and then extrapolated to the global terrestrial
ecosystems.

195 To evaluate the model's performance on capturing seasonal trends, the simulated N₂O emissions were plotted
along with observational data for the observational period. A linear regression is conducted to show the similarity
and discrepancy between simulations and observations, R², along with the slope and intercept of the linear
regression line are calculated. Six field observational sites having legume were organized for the model validation
(Table 3). Monthly N₂O emission data were used for model comparison.

2.5 Model Sensitivity and Uncertainty Analysis

200 The response of N fixation and N₂O emissions of different biomes to input data, and parameters was analyzed
using sensitivity test. Four major input variables were selected, including air temperature, precipitation, soil nitrogen
content and soil organic carbon content. The monthly average input variables were changed by ±10% of the original
level for each site and each grid. The variables were changed at 6 levels, respectively, and the rest of input variables
were kept at their original values. The sensitivity was calculated by comparing the simulated annual N₂O emissions
205 and nitrogen fixation to the simulations with the original input values (Figure 2).

3. Results

3.1 Model evaluation

To evaluate the model, thirty-five observational sites were selected for 7 major ecosystem types across the
globe, representing different climate and soil conditions. The experimental data of N fixation have a mean value of
210 12.9 kg N ha⁻¹ yr⁻¹, with a standard deviation of 17.7 kg N ha⁻¹ yr⁻¹. The maximum observed fixation occurred in
temperate forest in New Zealand, while the minimum rate was also for temperate forest in Idaho State of the US.
Our simulations are comparable with the observed data for all major ecosystem types with the coefficient of
determination (R²) of 0.44 and with a slope of 0.46 (Figure 3). The regression results are mainly influenced by some
observed data greater than 30 kg N ha⁻¹ yr⁻¹ (mean + standard deviation). By removing these data, the slope of
215 regression increases to 0.72. Observational data for temperate forests show the greatest variation among all major
ecosystem types, with a maximum value reaching 800 times of the minimum one. Simulations are closer to the
observations across sites in temperate forests with R² of 0.26 and slope of 0.42. Our model underestimated nitrogen
fixation rate in temperate forests. The large variation in observations may be due to the distribution of legume
plants, different sampling time periods (e.g., growing and non-growing seasons), and varying climate conditions.
220 For tropical forests, our model estimates of N fixation are higher than observations with the slope of 0.75 and R² of
0.44.



The model was also tested by comparing the simulated N₂O emissions with observed data for 5 sites and 3 ecosystem types including tropical forests, temperate forests, and grasslands. All observed data were converted into monthly average values. Our model reasonably reproduced the seasonal variations of N₂O emissions at
225 observational sites. In our earlier version of the model, the biological nitrogen input was assumed to be a constant throughout the year (Yu and Zhuang 2019). Compared to our earlier version, the current version contributes 5% to 20% of the total N₂O emission, but only leads to a minor difference to total seasonal trend throughout the year (Figure 3). There are still some discrepancies between simulated and observed N₂O emissions, which could be due to the uncertainty in measurement, sudden weather events and changes in soil characteristics. The comparison
230 between measured and simulated data further shows the influence of BNF for different ecosystem types on N₂O emissions. The influence is larger in tropical areas, while by percentage, BNF is more important for grasslands in temperate regions.

3.2 Model sensitivity analysis

The model sensitivity analysis quantifies the impact of changes in forcing data on nitrogen fixation rate and
235 N₂O emissions. Climate conditions including air temperature and precipitation, and soil characteristics of nitrogen content and carbon content varied at 3 levels to examine the sensitivity. The response of nitrogen fixation rate and N₂O emissions is quantified for each ecosystem type. The sensitivity test was conducted for all observational sites (Table 2 and Table 3). Temperature is the most sensitive variable (Figure 2). Comparing to N₂O emissions, nitrogen fixation is more sensitive to the change of all forcing conditions. Increasing soil nitrogen results in a lower N
240 fixation, but increasing N₂O emissions. Abundant soil nitrogen content inhibits BNF activity, but stimulates nitrification and denitrification processes.

3.3 Biological nitrogen fixation in global terrestrial ecosystems

Tropical forests in South America, Central Africa and South Asia show a wide range of N fixation rate between 1 and 200 kg N ha⁻¹ yr⁻¹ (Bruijnzeel et al, 1991). Here all plants in tropical rainforest are assumed to fix nitrogen and
245 one set of parameters are applied for all tropical forests. The coverage for tropical forests in the landscape was assumed to be 15% (Cleveland et al., 1999), ranging from 5% to 25%. The N₂ fixation rate was estimated to be 18.2 kg N ha⁻¹ yr⁻¹, which is the highest among all vegetation types. Our simulations show that the total fixed nitrogen ranges from 10.8 Tg N yr⁻¹ to 54 Tg N yr⁻¹, with the average value of 32.5 Tg N yr⁻¹ (Table 4). Nitrogen fixation in tropical forests is almost half of the global total amount and a principal contributor of BNF in natural ecosystems.
250 Tropical forests have the largest potential to fix nitrogen given that the optimal temperature and soil moisture for BNF is relatively easy to have under tropical climatic conditions.

Temperate forests cover the largest land area from 30°N to 60°N, including temperate coniferous forest, temperate deciduous forest and temperate evergreen forest. Temperate areas have the majority of legumes and many temperate ecosystems are considered to be N limited. Comparing to other ecosystem types in temperate regions,



255 conifers are likely to limit the reproduction of legumes (Wheatley et al, 2010). In general, plant species carrying
nitrogen fixers are only distributed in a small percentage of natural temperate forests, like clear-felled areas and
pastures (Boring and Swank, 1984). Cleveland et al. (1999) indicated that the legume coverage ranges from 1% to
10% of the land area only. Consequently, our simulations indicate that N fixation by temperate forests was 12.7 kg
N ha⁻¹ yr⁻¹. The estimates of the total nitrogen fixation were between 1.9 and 19.14 Tg N yr⁻¹ (Table 4). Nitrogen
260 fixation in temperate areas contributes 12.5% of the global total amount.

Savanna covers over a half of African continent, Australia and large areas of South America. It is an important
biome in the Southern Hemisphere. There is a great variation in native legume species. Only in humid savanna,
legumes may significantly contribute to the increase of soil nitrogen (Cech et al., 2008). On average, 15% of the
vegetation in savanna is considered as legume grass and biological nitrogen fixation occurs when precipitation is
265 greater than 10 mm per month. Generally, nitrogen fixation in savanna is restricted by soil moisture, while temperate
grassland is limited by both temperature and soil moisture (Bustamante et al, 1970). Nitrogen fixers are not abundant
for these biomes (Woodmansee et al., 1981). The coverage of nitrogen fixers was assumed to be from 5% to 25%,
(Cleveland et al., 1999). Our simulation assumed that nitrogen fixers cover 15% of the land, resulting in 1.9 kg N
ha⁻¹ yr⁻¹ fixation, representing a much smaller fraction compared to forest ecosystems. Total fixed nitrogen in
270 grasslands appeared to range from 0.62 to 3.1 Tg N yr⁻¹, with an average of 1.86 Tg N yr⁻¹. For savanna, the total
contribution was less due to its relatively small area. The minimum, average and maximum values were estimated to
be 0.45, 1.34 and 2.23 Tg N yr⁻¹, respectively.

In tundra and boreal forest regions, both host plants and their rhizobia are adapted to the environment with low
temperature. Nitrogen fixation rate is extremely variable for boreal ecosystems. For tundra, the coverage was
275 assumed to be 3-15%, and for boreal forest, the coverage was 4-18%. But in general, the low temperature and
permafrost conditions limit the activity of nitrogen fixers (Alexander, 1981). We estimated that tundra ecosystems
fix nitrogen at 3.2 kg N ha⁻¹ yr⁻¹. Their total BNF was between 0.51 to 2.55 Tg N yr⁻¹ with average of 1.54 Tg N yr⁻¹.
In boreal forests, the fixation rate was much lower (2.1 kg N ha⁻¹ yr⁻¹) compared to temperate forests.

In deserts, although some legumes may exist in extremely dry conditions, and some species may grow rapidly
280 after rainfall, their fixation could be neglected. However, in semi-arid areas, legumes are common plants with
several species, their N fixation is lower than tropical and temperate forests (5.7 kg N ha⁻¹ yr⁻¹).

Mediterranean ecosystems such as in south California and some areas in South Australia are characterized with
mild rainy winter and hot dry summer, containing both evergreen and deciduous shrublands, in which nodulated
legumes are prominent (Sprent et al., 2017). These legumes are more active in comparatively wet season than in dry
285 season (Sánchez-Díaz, 2001). The ability to fix nitrogen is considered to be one of the most important features that
enable legumes and plants to survive under severe environments (Crisp et al., 2004). We estimated that the N
fixation rate of these legume species is similar to that in grasslands (2.7 kg N ha⁻¹ yr⁻¹).

Spatially, the highest rate of N fixation occurred in the tropical and sub-tropical areas, as a result of proper
climate and soil characteristics for fixers (Figure 4a). N fixation from tropical forests and xeric shrubland contributes
290 to nearly half of the global terrestrial amount (Table 4). A lower N fixation rate was in high latitudes of East China,
North America and Europe, which were mainly covered with temperate forests. Compared to tropical areas, N



fixation in temperate regions shows a larger variability depending on vegetation types. The spatial variation could be attributed to the distribution of legume plants, in addition to the difference of humidity and temperature conditions. N fixation in temperate regions accounts for 35% of the total fixed N.

295 **3.4. Soil N₂O emissions enhanced by biological nitrogen fixation**

The model was run at site level to estimate the role of BNF on N₂O emissions. Our analysis between current model and previous model without N fixation module indicated that N₂O emissions would be affected by -5% to 20% depending on biome types and seasons (Figure 5). The involvement of BNF strengthens the seasonal variation as BNF led to lower emissions in winter and higher emissions in summer. Our model estimated that high BNF rates in growing season is consistent with other regional and global estimates (Cleveland et al., 1999, 2013; Lee and Son, 2005; Lett and Michelsen, 2014). The energetic cost for active N uptake becomes lowest when soil temperature is around 25°C (Fisher et al. 2010). Similarly, our estimates of high BNF rates also occur at similar temperature conditions in spring and summer. The global soil nitrogen mineralization rate was estimated to be 696 Tg N yr⁻¹ while 15% of plant N demand was provided by BNF (Cleveland et al., 2013). Our estimates of BNF were lower than the estimates by Cleveland et al. (2013) and fell within 10% of the total soil mineralization rate. This result also indicates that about 10% of the mineralized N was induced by BNF. As mineralized nitrogen is the substrate for subsequent reactions producing N₂O, the additional N input through BNF to terrestrial ecosystems led to the change of N₂O emissions across biomes.

During 1990-2000, our simulations show that BNF in natural terrestrial ecosystems is 61.5 Tg N yr⁻¹, but anthropogenic N fixation was much higher at 140 Tg N yr⁻¹ (Galloway et al., 2002). This large amount of anthropogenic N input to terrestrial ecosystems is expected to inhibit the natural BNF and might lead to less BNF in the future. Our simulations indicate that BNF has a minor influence on N₂O emissions for most high latitude ecosystems and extremely arid regions, as a result of low temperatures and drought (Figure 4b). Overall, the fixation as additional N to soils led to -5% to 20% changes in soil N₂O emissions.

315 **4. Discussion**

4.1 Comparison with other estimates of biological nitrogen fixation

There is a large uncertainty in estimating the N input into terrestrial ecosystems, especially from BNF (Sutton et al., 2014) (Table 4). In our study, a calibrated process-based model was applied to estimate site-level and global BNF in natural terrestrial ecosystems. Empirical models provide reasonable estimation based on relationships between fixation rate and environmental factors (e.g. evapotranspiration) (Cleveland et al, 1999), while process-based approaches consider processes in BNF affected by multiple controlling factors (Fisher et al., 2010; Gerber et al., 2008; Meyerholt et al., 2016). Our estimated BNF in the global terrestrial ecosystems is 61.5 Tg N yr⁻¹ with an uncertainty ranging from 19.8 to 107.9 Tg N yr⁻¹, which is lower than most existing studies. Cleveland et al. (1999) provided a central value of 195 Tg N yr⁻¹ by scaling up field-based experimental data, with a range of 100 - 289 Tg



325 N yr⁻¹. This range represents potential distribution of nitrogen fixation. In reality, N fixation is also affected by
climate and soil conditions, making the actual terrestrial BNF smaller than the potential one. In a more recent study
of Cleveland et al. (2013), a total of 127.5 Tg N yr⁻¹ was estimated to be related to BNF, based on the relationship
between BNF and evapotranspiration (ET). Galloway et al (2002b) also provided several estimates for global BNF.
Galloway et al. (2004) further suggested a range of 100 - 290 Tg N yr⁻¹ and implied that the true rate of BNF would
330 be at the low end of this range without large-scale human disturbance. In an earlier study (Galloway et al., 2002b),
the mean annual global BNF was estimated to be 89-100 Tg N yr⁻¹. By assuming a steady state between N input to
and loss from ecosystems, Vitousek et al. (2013) estimated the BNF to be 58 Tg N yr⁻¹ with a plausible range of 40 -
100 Tg N yr⁻¹, which is similar to our estimates. However, Xu-Ri and Prentice (2017) estimated that the N fixation
was about 340 Tg N yr⁻¹ which is almost 5 times larger than our estimates. In their study, BNF was determined by
335 plant N requirement across all biome types.

In our estimation, tropical forests significantly contribute to the total BNF, which is up to 18 kg N ha⁻¹yr⁻¹. This
result is highly related to the density of leguminous plants, and the physical conditions in tropical areas (Crews,
1999). Our simulated results are comparable to the estimates of symbiotic N₂ fixation from tropical evergreen (5.5-
16 kg N ha⁻¹ yr⁻¹) and deciduous forests (7.5-30 kg N ha⁻¹ yr⁻¹) (Reed et al., 2011). Barron et al. (2010) directly
340 measured N₂-fixing root nodules across lowland tropical forests and their observations also showed a large variation
among individual trees. For a mature forest matrix, the average value was around 10 kg N ha⁻¹ yr⁻¹, but it could be as
high as 200 kg N ha⁻¹ yr⁻¹ for some areas. Cleveland (2013) provided a similar estimate to ours (around 12 kg N ha⁻¹
yr⁻¹), but higher values (20-30 kg N ha⁻¹ yr⁻¹) in their earlier studies (Cleveland et al., 1999). Sullivan et al (2014)
analyzed human's impact on tropical N fixation and found, depending on forest ages, fixation was 5.7 kg N ha⁻¹ yr⁻¹
345 with a range from 1.2 to 14.4 kg N ha⁻¹ yr⁻¹, which is lower than our estimates.

For temperate and boreal forests, we estimated that BNF fixation is 2.1-18 kg N ha⁻¹ yr⁻¹. The existing BNF
estimates from literature also show a large uncertainty for those forest ecosystems. For instance, LM3V-N model
(Gerber et al., 2009) suggested that the N input to forests to be less than 5 kg N ha⁻¹ yr⁻¹. But their model also
estimated that, in moist forests, the uptake of N could be 30-80 kg N ha⁻¹ yr⁻¹. Deluca et al. (2002) reported that
350 cyanobacterium and feather moss could act as a supplement to N fixation in boreal forests (0.5 kg N ha⁻¹ yr⁻¹) while
the organic N accumulation could be 3 kg N ha⁻¹ yr⁻¹. For the forests in northwest Rocky Mountain, N fixation
amount is on average between 0.5 and 2 kg N ha⁻¹ yr⁻¹ (Clayton and Kennedy, 1985; Fahey et al., 1988) while Kou-
Giesbrecht and Menge's model (2019) estimated the N fixation rate to be 0 -10 kg N ha⁻¹ yr⁻¹ for temperate forests,
and 0 to 6 kg N ha⁻¹ yr⁻¹ for boreal forests.

355 There could be a number of reasons for our comparatively lower estimates. The most important one is that there
is a considerable uncertainty in estimating the coverage of N fixing plants. High diversity in the distribution of
legume plants highly influences the estimation of total plant coverage, because our estimation was based on site-
level experimental data. In order to improve our understanding, more investigation on legume plant distribution and
associated data for N fixers is needed, especially in the Middle Asia, South America and Africa.

360 Large variations of BNF rates exist across terrestrial ecosystems spatially (Figure 4a). The global BNF spatial
pattern is similar to other estimates (Cleveland et al., 1999; Xu-Ri and Prentice, 2017). The highest N fixation rate in



365

tropical regions (more than 50% of the global terrestrial N fixation) is primarily due to their warm and moist soil conditions. Further, N fixed by human activities became increasingly influential in the past century (Galloway et al., 2002), especially in temperate regions due to their large human population. The anthropogenic N deposition contributed more to soil N than BNF did. As a result, soils became N rich, inhibiting BNF in temperate soils. This could explain why the potential N fixation rate was high in temperate ecosystems, but only contributed to 20% of the total fixation.

4.2 Major controls to biological nitrogen fixation

370

In our simulations, the N fixation was primarily influenced by soil temperature, moisture and soil nitrogen content. The highest N fixation rate in tropical ecosystems is consistent with our sensitivity analysis for temperature and soil moisture. The sensitivity analysis indicated that a 1-3°C increase of temperature led to 7% increase in N fixation rate. Nitrogen cycle responds differently between different biomes and legume types. But in general, increasing temperature will accelerate processes in the N cycle. Soil moisture correlates with BNF in a similar way with temperature. A slightly increase of precipitation (10%) increased the nitrogenase activity. However, the response of N fixation to soil water stress is not as sensitive as that to the change in temperature. Xeric shrubland and savanna in dry tropical areas still contribute greatly to the global N fixation, while the contribution of boreal forests, with abundant rainforest and low temperature, is much lower.

375

380

BNF is highly regulated by soil nitrogen content. N-deficiency conditions usually favor BNF activities, for example, in xeric shrubland and savanna. Enhancing soil N content will decrease the N fixation rate, which is also consistent with our sensitivity analysis. It costs less energy for plants to take up N directly from soils rather than biologically fixing it from the atmosphere (Cannell and Thornley, 2000). However, there is an exception for some areas in tropical ecosystems. Many tropical soils are comparatively rich in nitrogen, but N-fixing plants are still active to compensate the nitrogen depletion due to the rapid N cycling (Pons et al., 2007). This explains why N fertilization inhibits the BNF in temperate ecosystems, but BNF is still active in N-rich soils in tropical ecosystems. In areas where the energetic cost succeeds the demand of N, the BNF rate will be comparatively lower. Sullivan et al. (2014) suggested that there were lower rates of BNF in undisturbed mature forests and higher rate in secondary forests, depending on the balance between N-demand and energy consumption.

385

4.3 Model limitation and future work

390

The incorporation of BNF into TEM allows us to more adequately estimate N₂O emissions from natural terrestrial ecosystems. However, there are several limitations in this study.

First, the current model ignores the effect of free-living BNF. Although symbiotic BNF is critical for most natural and semi-natural ecosystems, asymbiotic organisms play an important role in extreme environments such as waterlogged soils and deserts. The importance of symbiotic BNF or fixation by leguminous plants may not be as significant as previously thought. Elbert et al. (2012) suggested that cryptogam contributed nearly half BNF in



395 terrestrial ecosystems, which was up to 49 Tg N yr^{-1} . In some tropical areas, the spatial N input from free-living
bacteria even exceeds symbiotic input (Sullivan et al., 2014). In addition, legumes are not the only source of
symbiotic BNF. Some fungi species have the ability to actively fix atmospheric nitrogen. But in most existing
models, fungi or mycorrhizae symbioses are not considered due to the limited knowledge about their mechanisms of
fixing N (Fisher et al., 2010). A more comprehensive model that covers various types of nitrogen fixation is needed.

400 Second, the BNF process in our model is calibrated with a limited amount of data, imposing a general set of
parameters to all plant species and soil conditions within an ecosystem type. More observational data from natural
terrestrial ecosystems is desirable to improve our model.

Third, it is difficult to isolate the N addition via natural processes from human activities. In the US, 20-35% of
annual N input into terrestrial ecosystems are human-related (Sobata et al., 2013). As a result, the quality of
405 observational data varies from site to site, some BNF data are only semi-natural. The observational data are
imperfect, which might have also biased our estimates through model parameterization process.

5. Conclusions

This study developed a process-based biological nitrogen fixation model and coupled it with an extant nitrous
oxide emission model. The model was evaluated with observed data for both N fixation and N_2O emissions. The
410 model was then extrapolated to the global natural terrestrial ecosystems. Our model estimated that biological
nitrogen fixation in natural terrestrial ecosystems was $61.5 \text{ Tg N yr}^{-1}$ during the last decade of the 20th century and
the greatest fixation rate occurred in tropical regions. Soil temperature, rather than soil moisture and nutrient
content, is the most dominant control to the fixation. We further found that, the fixation as additional N to soils led
to -5% to 20% changes in soil N_2O emissions. Both the new and previous models were able to capture the seasonal
415 variation of field-observed N_2O , but the new version of the model estimated more seasonal dynamics. Lacking the
knowledge about the distribution of N fixing plants and their physiological features might have biased our estimates
of both biological nitrogen fixation and N_2O emissions at the global scale.

Data availability

420 Climate data including monthly cloudiness, precipitation, temperature, and water vapor pressure are from the
Climate Research Unit (CRU) <http://www.cru.uea.ac.uk/data> (last access: May 2017). Global vegetation data and
soil data are available in Zhuang et al. (2003) and McGuire et al. (2001). The explicit spatial data on soil water pH
from the ORDL gridded soil properties product (https://daac.ornl.gov/cgi-bin/dsvviewer.pl?ds_id=546, last access:
March ,2017) are based on the World Inventory of Soil Emission Potentials (WISE) database (Batjes, 2000). The
425 global average carbon dioxide concentration is observed at NOAA's Mauna Loa Observatory. N deposit data are
from NADP monitor and CASTNET. The initial values of soil microbial carbon and nitrogen, and the ratio of C / V /
N at the global scale, were from a compilation of global soil microbial biomass carbon, nitrogen, and phosphorus
data (<https://doi.org/10.3334/ORNLDAAAC/1264>, last access: May, 2017). *The data presented in this paper can be
accessed through our research website (<http://www.eaps.purdue.edu/ebdl/>).*

430



Author contribution

Q. Zhuang and T. Yu designed the research. T. Yu performed model simulations and data analysis. Both authors contributed to the paper writing.

435 **Competing interests**

The authors declare that they have no conflict of interest.

Acknowledgments

440 This study is supported through projects funded by the NASA Land Use and Land Cover Change program (NASA-NNX09AI26G), Department of Energy (DE-FG02-08ER64599), the NSF Division of Information & Intelligent Systems (NSF-1028291). Thanks to Rosen Center for Advanced Computing (RCAC) at Purdue University for computing support.



445 Reference:

- Adams M A, Attiwill P M. Role of *Acacia* spp. in nutrient balance and cycling in regenerating *Eucalyptus regnans* F. Muell. forests. I. Temporal changes in biomass and nutrient content. *Aust. J. Bot.*, 32(2): 205-215, 1984.
- AGAGE (Advanced Global Atmospheric Gases Experiment). 2015. Monthly mean N₂O concentrations for Cape Grim, Australia. Accessed June 5, 2015.
- 450 Anderson I C, Poth M A. Semiannual losses of nitrogen as NO and N₂O from unburned and burned chaparral. *Global Biogeochem. Cy.*, 3(2): 121-135, 1989.
- Alexander V, Billington M M. Nitrogen fixation in the Alaskan taiga, *Forest ecosystems in the Alaskan taiga*. Springer, New York, NY, 112-120, 1986.
- Baker T G, Oliver G R, Hodgkiss P D. Distribution and cycling of nutrients in *Pinus radiata* as affected by past lupin growth and fertiliser. *Forest Ecol. Manag.*, 17(2-3): 169-187, 1986.
- 455 Barron A R, Purves D W, Hedin L O. Facultative nitrogen fixation by canopy legumes in a lowland tropical forest. *Oecologia*, 165(2): 511-520, 2011.
- Battle M, Bender M, Sowers T, et al. Atmospheric gas concentrations over the past century measured in air from firn at the South Pole. *Nature*, 383(6597): 231, 1996.
- 460 Beck, Douglas, and Luis A. Materon, eds. *Nitrogen Fixation by Legumes in Mediterranean Agriculture: Proceedings of a Workshop on Biological Nitrogen Fixation on Mediterranean-type Agriculture*, ICARDA, Syria, April 14–17, 1986. Vol. 32. Springer Science & Business Media, 2012.
- Blundon D J, Dale M R T. Dinitrogen fixation (acetylene reduction) in primary succession near Mount Robson, British Columbia, Canada[J]. *Arctic. Alpine. Res.*, 22(3): 255-263, 1990.
- 465 Boote K J, Jones J W, Hoogenboom G, et al. Simulation of crop growth: CROPGRO model. *Agricultural systems modeling and simulation*, 18: 651-692, 1998.
- Boring L R, Swank W T, Waide J B, et al. Sources, fates, and impacts of nitrogen inputs to terrestrial ecosystems: review and synthesis. *Biogeochemistry*, 6(2): 119-159, 1988.
- Boring L R, Swank W T. The role of black locust (*Robinia pseudo-acacia*) in forest succession. *J. Ecol.*, 749-766, 470 1984.
- Bowman W D, Schardt J C, Schmidt S K. Symbiotic N₂-fixation in alpine tundra: ecosystem input and variation in fixation rates among communities. *Oecologia*, 108(2): 345-350, 1996.
- Breitbarth E, Oschlies A, LaRoche J. Physiological constraints on the global distribution of *Trichodesmium*? effect of temperature on diazotrophy[J]. *Biogeosciences*, 4(1): 53-61, 2007.
- 475 Bruijnzeel L A. Nutrient input–output budgets of tropical forest ecosystems: a review. *J. Trop. Ecol.*, 7(1): 1-24, 1991.
- Bustamante, M. et al. Nitrogen cycling in tropical and temperate savannas. 10.1007/978-1-4020-5517-1_10, 1970.
- Cannell M G R, Thornley J H M. Modelling the components of plant respiration: some guiding principles. *Ann. Bot. London*, 85(1): 45-54, 2000.
- Cech P G, Kuster T, Edwards P J, et al. Effects of herbivory, fire and N₂-fixation on nutrient limitation in a humid 480 African savanna. *Ecosystems*, 11(6): 991, 2008.



- Chen G X, Huang B, Xu H, et al. Nitrous oxide emissions from terrestrial ecosystems in China. *Chemosphere-Global Change Science*, 2(3-4): 373-378, 2000.
- Chen M I N, Zhuang Q. Modelling temperature acclimation effects on the carbon dynamics of forest ecosystems in the conterminous United States. *Tellus B.*, 65(1): 1915, 2013.
- 485 Clayton J L, Kennedy D A. Nutrient Losses from Timber Harvest in the Idaho Batholith I. *Soil. Sci. Soc. Am. J.*, 49(4): 1041-1049, 1985.
- Cleveland C C, Townsend A R, Schimel D S, et al. Global patterns of terrestrial biological nitrogen (N₂) fixation in natural ecosystems. *Global biogeochem. Cy.*, 13(2): 623-645, 1999.
- Cleveland C C, Houlton B Z, Smith W K, et al. Patterns of new versus recycled primary production in the terrestrial biosphere. *P. Natl. A. Sci.*, 110(31): 12733-12737, 2013.
- 490 Corre-Hellou G, Brisson N, Launay M, et al. Effect of root depth penetration on soil nitrogen competitive interactions and dry matter production in pea–barley intercrops given different soil nitrogen supplies. *Field Crop. Res.*, 103(1): 76-85, 2007.
- Corre-Hellou G, Faure M, Launay M, et al. Adaptation of the STICS intercrop model to simulate crop growth and N accumulation in pea–barley intercrops. *Field Crop. Res.*, 113(1): 72-81, 2009.
- 495 Crews T E. The presence of nitrogen fixing legumes in terrestrial communities: Evolutionary vs ecological considerations, *New Perspectives on Nitrogen Cycling in the Temperate and Tropical Americas*. Springer, Dordrecht, 233-246, 1999.
- Crisp M, Cook L, Steane D. Radiation of the Australian flora: what can comparisons of molecular phylogenies across multiple taxa tell us about the evolution of diversity in present–day communities?. *Philos. T. Roy. Soc. B.*, 500 359(1450): 1551-1571, 2004.
- Davidson E A, Reis de Carvalho C J, Vieira I C G, et al. Nitrogen and phosphorus limitation of biomass growth in a tropical secondary forest. *Ecol. Appl.*, 14(sp4): 150-163, 2004.
- Dentener F, Drevet J, Lamarque J F, et al. Nitrogen and sulfur deposition on regional and global scales: A multimodel evaluation. *Global biogeochemical cycles*, 20(4), 2006.
- 505 DuBois J D, Kapustka L A. Biological nitrogen influx in an Ohio relict prairie. *American Journal of Botany*, 70(1): 8-16, 1983.
- Duxbury, J. M., Bouldin, D. R., Terry, R. E., & Tate Iii, R. L.. Emissions of nitrous oxide from soils. *Nature*, 298(5873), 462, 1982.
- 510 Eckersten, H., Geijersstam, L. A., & Torssell, B. Modelling nitrogen fixation of pea (*Pisum sativum* L.). *Acta. Agr. Scand. B-S P*, 56(2), 129-137, 2006.
- Eisele, L., Schimel, D. S., Kapustka, L. A., & Parton, W. J. Effects of available P and N: P ratios on non-symbiotic dinitrogen fixation in tallgrass prairie soils. *Oecologia*, 79(4), 471-474, 1989.
- Elbert, W., Weber, B., Burrows, S., Steinkamp, J., Büdel, B., Andreae, M. O., & Pöschl, U. Contribution of cryptogamic covers to the global cycles of carbon and nitrogen. *Nat. Geosci.*, 5(7), 459, 2012.
- 515 Erickson H, Davidson E A, Keller M. Former land-use and tree species affect nitrogen oxide emissions from a tropical dry forest. *Oecologia*, 130(2): 297-308, 2002.



- Erickson H, Keller M, Davidson E A. Nitrogen oxide fluxes and nitrogen cycling during postagricultural succession and forest fertilization in the humid tropics. *Ecosystems*, 4(1): 67-84, 2001.
- 520 Fahey T J, Yavitt J B, Pearson J A, et al. The nitrogen cycle in lodgepole pine forests, southeastern Wyoming. *Biogeochemistry*, 1(3): 257-275, 1985.
- Fisher J B, Sitch S, Malhi Y, et al. Carbon cost of plant nitrogen acquisition: A mechanistic, globally applicable model of plant nitrogen uptake, retranslocation, and fixation. *Global Biogeochem. Cy.*, 24(1), 2010.
- Galloway J N, Schlesinger W H, Levy H, et al. Nitrogen fixation: Anthropogenic enhancement-environmental
525 response. *Global Biogeochem. Cy.*, 9(2): 235-252, 1995.
- Galloway J N, Cowling E B, Seitzinger S P, et al. Reactive nitrogen: too much of a good thing?. *AMBIO: A Journal of the Human Environment*, 31(2): 60-64, 2002.
- Galloway, James N., and Ellis B. Cowling. "Reactive nitrogen and the world: 200 years of change." *AMBIO: A Journal of the Human Environment* 31.2 (2002): 64-71.
- 530 Galloway J N, Dentener F J, Capone D G, et al. Nitrogen cycles: past, present, and future. *Biogeochemistry*, 70(2): 153-226, 2004.
- Gerber S, Hedin L O, Oppenheimer M, et al. Nitrogen cycling and feedbacks in a global dynamic land model. *Global Biogeochem. Cy.* 24(1), 2010.
- Goossens A, De Visscher A, Boeckx P, et al. Two-year field study on the emission of N₂O from coarse and middle-textured Belgian soils with different land use. *Nutr Cycl Agroecosys.*, 60(1-3): 23-34, 2001.
- 535 Gregory J, Stouffer R J, Molina M, et al. *Climate change 2007: the physical science basis*. 2007.
- Grove T S, Malajczuk N. Nodule production and nitrogen fixation (acetylene reduction) by an understorey legume (*Bossiaea laidlawiana*) in Eucalyptus forest. *J. Ecol.*, 303-314, 1992.
- Gundale M J, Nilsson M, Bansal S, et al. The interactive effects of temperature and light on biological nitrogen fixation
540 in boreal forests. *New Phytol*, 194(2): 453-463, 2012.
- Hardy R W F, Holsten R D, Jackson E K, et al. The acetylene-ethylene assay for N₂ fixation: laboratory and field evaluation. *Plant physiology*, 1968, 43(8): 1185-1207, 1968.
- Hardy R W F, Burns R C, Holsten R D. Applications of the acetylene-ethylene assay for measurement of nitrogen fixation. *Soil Biol. Biochem.*, 5(1): 47-81, 1973.
- 545 Harvey A E, Larsen M J, Jurgensen M F, et al. Nitrogenase activity associated with decayed wood of living northern Idaho conifers. *Mycologia*, 81(5): 765-771, 1989.
- He X H, Critchley C, Bledsoe C. Nitrogen transfer within and between plants through common mycorrhizal networks (CMNs). *Crit Rev Plant. Sci*, 22(6): 531-567, 2003.
- Heath, B., Sollins, P., Perry, D. A., & Cromack Jr, K. Asymbiotic nitrogen fixation in litter from Pacific Northwest
550 forests. *Can J. Forest Res.*, 18(1), 68-74, 1988.
- Hendrickson, O. Q. Asymbiotic nitrogen fixation and soil metabolism in three Ontario forests. *Soil Biol. Biochem.*, 22(7), 967-971, 1990.



- 555 Holzworth, D. P., Huth, N. I., deVoil, P. G., Zurcher, E. J., Herrmann, N. I., McLean, G., ... & Moore, A. D. APSIM–
evolution towards a new generation of agricultural systems simulation. *Environ. Modell. Softw.*, 62, 327-350,
2014.
- Houlton, B. Z., Morford, S. L., & Dahlgren, R. A. Convergent evidence for widespread rock nitrogen sources in
Earth's surface environment. *Science*, 360(6384), 58-62, 2018.
- Huss-Danell, K. Nitrogen fixation by *Stereocaulon paschale* under field conditions. *Can J. Botany*, 55(5), 585-592,
1977.
- 560 Jarrell, W. M., & Virginia, R. A. Soil cation accumulation in a mesquite woodland: sustained production and long-
term estimates of water use and nitrogen fixation. *J. Arid Environ.* 18(1), 51-58, 1990.
- Johnson, H. B., & Mayeux, H. S. *Prosopis glandulosa* and the nitrogen balance of rangelands: extent and occurrence
of nodulation. *Oecologia*, 84(2), 176-185, 1990.
- Kapustka, L. A., & DuBois, J. D. Dinitrogen fixation by cyanobacteria and associative rhizosphere bacteria in the
565 Arapaho Prairie in the Sand Hills of Nebraska. *Am. J. Bot.*, 74(1), 107-113, 1987.
- Lambers, H., Raven, J. A., Shaver, G. R., & Smith, S. E. Plant nutrient-acquisition strategies change with soil age.
Trends Ecol. Evol., 23(2), 95-103, 2008.
- LeBauer, D. S., & Treseder, K. K. Nitrogen limitation of net primary productivity in terrestrial ecosystems is globally
distributed. *Ecology*, 89(2), 371-379, 2008.
- 570 Lee, Y. Y., & Son, Y. Diurnal and seasonal patterns of nitrogen fixation in *Analus hirsuta* plantation of central Korea.
J. Plant Biol., 48(3), 332-337, 2005.
- Lepper, M. G., & Fleschner, M. Nitrogen fixation by *Cercocarpus ledifolius* (Rosaceae) in pioneer habitats. *Oecologia*,
27(4), 333-338, 1977.
- Lett, S., & Michelsen, A. Seasonal variation in nitrogen fixation and effects of climate change in a subarctic heath.
575 *Plant Soil*, 379(1-2), 193-204, 2014.
- Levy, H., Moxim, W. J., & Kasibhatla, P. S. A global three-dimensional time-dependent lightning source of
tropospheric NO_x. *J Geophys. Res-Atmos.*, 101(D17), 22911-22922, 1996.
- Li, C., Aber, J., Stange, F., Butterbach-Bahl, K., & Papen, H. A process-oriented model of N₂O and NO emissions
from forest soils: 1. Model development. *J Geophys. Res-Atmos.*, 105(D4), 4369-4384, 2000.
- 580 Lindemann, William C., and C. R. Glover. Nitrogen fixation by legumes, 2003.
- Maheswaran, J., & Gunatilleke, I. A. U. N. Nitrogenase activity in soil and litter of a tropical lowland rain forest and
an adjacent fernland in Sri Lanka. *J Trop Ecol*, 6(3), 281-289, 1990.
- Marino, D., Frendo, P., Ladrera, R., Zabalza, A., Puppo, A., Arrese-Igor, C., & González, E. M. Nitrogen fixation
control under drought stress. Localized or systemic?. *Plant Physiol.*, 143(4), 1968-1974, 2007.
- 585 Matson, P. A., Gower, S. T., Volkman, C., Billow, C., & Grier, C. C. Soil nitrogen cycling and nitrous oxide flux in
a Rocky Mountain Douglas-fir forest: effects of fertilization, irrigation and carbon addition. *Biogeochemistry*,
18(2), 101-117, 1992.
- Matson, P. A., Volkman, C., Coppinger, K., & Reiners, W. A. Annual nitrous oxide flux and soil nitrogen
characteristics in sagebrush steppe ecosystems. *Biogeochemistry*, 14(1), 1-12, 1991.



- 590 May, D.E., and Webber, P.J., Spatial and temporal variation of vegetation and its productivity on Niwot Ridge,
Colorado, in *Ecological Studies in the Colorado Alpine*, a Festschrift for John W. Mart, edited by H.
Halfpenny, pp. 35-62, Institute for Arctic and Alpine Research, Univ. of Colorado, Boulder, Colo., 1982.
- Mitchell, T. D., & Jones, P. D. An improved method of constructing a database of monthly climate observations and
associated high-resolution grids. *Int J Climatol.: A Journal of the Royal Meteorological Society*, 25(6), 693-
595 712, 2005.
- Montanez, A., Danso, S. K. A., & Hardarson, G. (1995). The effect of temperature on nodulation and nitrogen fixation
by five *Bradyrhizobium japonicum* strains. *Appl. Soil Ecol.*, 2(3), 165-174.
- Morford, S. L., Houlton, B. Z., & Dahlgren, R. A. Increased forest ecosystem carbon and nitrogen storage from
nitrogen rich bedrock. *Nature*, 477(7362), 78, 2011.
- 600 Mosier, A. R., Delgado, J. A., Cochran, V. L., Valentine, D. W., & Parton, W. J. Impact of agriculture on soil
consumption of atmospheric CH₄ and a comparison of CH₄ and N₂O flux in subarctic, temperate and
tropical grasslands. *Nutr. Cycl. Agroecosys.*, 49(1-3), 71-83, 1997.
- Mosier, A. R., Parton, W. J., Valentine, D. W., Ojima, D. S., Schimel, D. S., & Heinemeyer, O. CH₄ and N₂O fluxes
in the Colorado shortgrass steppe: 2. Long-term impact of land use change. *Global Biogeochem. Cy.*, 11(1),
605 29-42, 1997..
- Mus, F., Crook, M. B., Garcia, K., Costas, A. G., Geddes, B. A., Kouri, E. D., ... & Udvardi, M. K. Symbiotic nitrogen
fixation and the challenges to its extension to nonlegumes. *Appl. Environ. Microbiol.*, 82(13), 3698-3710,
2016.
- Müller, C., & Sherlock, R. R.. Nitrous oxide emissions from temperate grassland ecosystems in the Northern and
610 Southern Hemispheres. *Global Biogeochem. Cy.*, 18(1), 2004.
- NOAA (National Oceanic and Atmospheric Administration). 2016. Monthly mean N₂O concentrations for Barrow,
Alaska; Mauna Loa, Hawaii; and the South Pole. Accessed June 8, 2016.
- O'Connell, A. M., & Grove, T. S. Seasonal variation in C₂H₂ reduction (N₂-fixation) in the litter layer of eucalypt
forests of south-western Australia. *Soil Biol Biochem.*, 19(2), 135-142, 1987.
- 615 Omari, K., Mubyana, T., Matsheka, M. I., Bonyongo, M. C., Veenendaal, E., & Musil, C. F. Flooding and its influence
on diazotroph populations and soil nitrogen levels in the Okavango Delta. *S Afr. J. Bot.*, 70(5), 734-740,
2004.
- Permar, T. A., & Fisher, R. F. Nitrogen fixation and accretion by wax myrtle (*Myrica cerifera*) in slash pine (*Pinus
elliottii*) plantations. *Forest Ecol Manag.*, 5(1), 39-46, 1983.
- 620 Pitcairn, C. E. R., Skiba, U. M., Sutton, M. A., Fowler, D., Munro, R., & Kennedy, V. Defining the spatial impacts of
poultry farm ammonia emissions on species composition of adjacent woodland groundflora using Ellenberg
Nitrogen Index, nitrous oxide and nitric oxide emissions and foliar nitrogen as marker variables. *Environ.
Po.*, 119(1), 9-21, 2002.
- Pons, T. L., Perreijn, K., Van Kessel, C., & Werger, M. J. Symbiotic nitrogen fixation in a tropical rainforest: 15N
625 natural abundance measurements supported by experimental isotopic enrichment. *New Phytol.*, 173(1), 154-
167, 2007.



- Prentice, I. C. Modelling the demand for new nitrogen fixation by terrestrial ecosystems. *Biogeosciences*, 14(7), 2003-2017, 2017.
- 630 Reed, S. C., Cleveland, C. C., & Townsend, A. R. Functional ecology of free-living nitrogen fixation: a contemporary perspective. *Annu Rev Ecol. Evol. S.*, 42, 489-512, 2011.
- Rosswall T, Granhall U. Nitrogen cycling in a subarctic ombrotrophic mire. *Ecol. Bull.*: 209-234, 1980.
- Rundel, P. W., Nilsen, E. T., Sharifi, M. R., Virginia, R. A., Jarrell, W. M., Kohl, D. H., & Shearer, G. B.. Seasonal dynamics of nitrogen cycling for a *Prosopis* woodland in the Sonoran Desert. In *Nitrogen Cycling in Ecosystems of Latin America and the Caribbean* (pp. 343-353), 1982. Springer, Dordrecht..
- 635 Sánchez-Díaz, M. Adaptation of legumes to multiple stresses in Mediterranean-type environments. *Options Méditerranéennes*, 45, 145-151, 2001.
- Schlesinger, W. H., Gray, J. T., Gill, D. S., & Mahall, B. E. *Ceanothus megacarpus* chaparral: a synthesis of ecosystem processes during development and annual growth. *Bot. Rev.*, 48(1), 71-117, 1982.
- Schlesinger, W. H. Biogeochemical Cycles.(Book Reviews: Biogeochemistry. An Analysis of Global Change.).
640 *Science*, 253, 686-687, 1991.
- Schlesinger, W. H. On the fate of anthropogenic nitrogen. *P Natl. Acad.*, 106(1), 203-208, 2009.
- Schrire, B. D., G. P. Lewis, and M. Lavin. Biogeography of the Leguminosae. *Legumes of the world*: 21-54, 2005..
- Schwintzer, C. R. Nonsymbiotic and symbiotic nitrogen fixation in a weakly minerotrophic peatland. *Am. J. Bot.*, 70(7), 1071-1078., 1983.
- 645 Shearer, G., & Kohl, D. H. N₂-fixation in field settings: estimations based on natural ¹⁵N abundance. *Funct. Plant Biol.*, 13(6), 699-756, 1986.
- Sheridan, R. P. Nitrogenase activity by *Hapalosiphon flexuosus* associated with *Sphagnum erythrocalyx* mats in the cloud forest on the volcano La Soufriere, Guadeloupe, French West Indies. *Biotropica*, 134-140, 1991.
- Skiba, U. M., Sheppard, L. J., MacDonald, J., & Fowler, D. Some key environmental variables controlling nitrous oxide emissions from agricultural and semi-natural soils in Scotland. *Atmos. Environ.*, 32(19), 3311-3320, 1998.
- 650 Skujiņš, J., Tann, C. C., & Börjesson, I. Dinitrogen fixation in a montane forest sere determined by ¹⁵N₂ assimilation and in situ acetylene-reduction methods. *Soil Biol. Biochem.*, 19(4), 465-471, 1987.
- Sobota, D. J., Compton, J. E., & Harrison, J. A. Reactive nitrogen inputs to US lands and waterways: how certain are we about sources and fluxes?. *Front. Ecol. Environ.*, 11(2), 82-90, 2013.
- 655 Sonesson, M., Jonsson, S., Rosswall, T., & Rydén, B. E. The Swedish IBP/PT Tundra Biome Project Objectives-Planning-Site. *Ecol. Bull.*, 7-25, 1980.
- Sprent, J. I. The effects of water stress on nitrogen-fixing root nodules. *New Phytol.*, 71(3), 443-450, 1972.
- Sprent, J. I., Ardley, J., & James, E. K. Biogeography of nodulated legumes and their nitrogen-fixing symbionts. *New
660 Phytol.*, 215(1), 40-56, 2017.
- Srivastava, A. K., & Ambasht, R. S. Soil moisture control of nitrogen fixation activity in dry tropical *Casuarina* plantation forest. *J. Environ. Manage.*, 42(1), 49-54, 1994.



- Stedman, D. H., and R. Shetter, The global budget of atmosphere nitrogen species, in Trace Atmospheric Constituent: Properties, Transformations and Fates, edited by S.S. Schwartz, pp. 411-454, John Wiley, New York, 1983.
- 665 Sutton, Mark A., et al., eds. Nitrogen deposition, critical loads and biodiversity. Springer Science & Business Media, 2014.
- Sullivan, B. W., Smith, W. K., Townsend, A. R., Nasto, M. K., Reed, S. C., Chazdon, R. L., & Cleveland, C. C. (2014). Spatially robust estimates of biological nitrogen (N) fixation imply substantial human alteration of the tropical N cycle. *P. Natl. A. Sci.*, 111(22), 8101-8106, 2014.
- 670 Thornley, J. H. M. Simulating grass-legume dynamics: a phenomenological submodel. *Ann Bot-London*, 88(5), 905-913, 2001.
- Veldkamp, E., Keller, M., & Nuñez, M. Effects of pasture management on N₂O and NO emissions from soils in the humid tropics of Costa Rica. *Global Biogeochem. Cy.*, 12(1), 71-79, 1998.
- Verchot, L. V., Hutabarat, L., Hairiah, K., & Van Noordwijk, M. Nitrogen availability and soil N₂O emissions following conversion of forests to coffee in southern Sumatra. *Global Biogeochem. Cy.*, 20(4), 2006.
- 675 Vitousek, P. M. Potential nitrogen fixation during primary succession in Hawaii Volcanoes National Park. *Biotropica*, 234-240, 1994.
- Vitousek, P. M., Menge, D. N., Reed, S. C., & Cleveland, C. C. Biological nitrogen fixation: rates, patterns and ecological controls in terrestrial ecosystems. *Philos. T. R. Soc. B.*, 368(1621), 20130119, 2013.
- 680 Voisin, A. S., Salon, C., Jeudy, C., & Warembourg, F. R. Symbiotic N₂ fixation activity in relation to C economy of *Pisum sativum* L. as a function of plant phenology. *J. Exp. Bot.*, 54(393), 2733-2744, 2003.
- Voisin, A. S., Bourion, V., Duc, G., & Salon, C. Using an ecophysiological analysis to dissect genetic variability and to propose an ideotype for nitrogen nutrition in pea. *Ann Bot-London*, 100(7), 1525-1536, 2007.
- Wheatley, Ron E., and J. I. Sprent. Legume Nodulation: A Global Perspective. *Exp. Agr.* 46.4: 568, 2010.
- 685 Whitehead, D.C., Grass Nitrogen, 397pp., CAB Int., Wallingford, UK
- Williams, J. R. Sharply A N. EPIC-Erosion Productivity Impact Calculator | . Model Documentation. US Department of Agriculture Technical Bulletin, (1990), 1768.
- Woodmansee, R. G., & Wallach, L. S. Effects of fire regimes on biogeochemical cycles. *Terrestrial Nitrogen Cycles. Ecol. Bull.(Stockholm)*, 33, 649-669, 1981.
- 690 Wu, L., & McGechan, M. B. Simulation of nitrogen uptake, fixation and leaching in a grass/white clover mixture. *Grass Forage Sci.*, 54(1), 30-41, 1999.
- Yu, T. and Q. Zhuang. Quantifying global N₂O emissions from natural ecosystem soils using trait-based biogeochemistry models. *Biogeosciences* 16(2): 207-222, 2019.
- Zhuang, Q., Romanovsky, V. E., & McGuire, A. D. Incorporation of a permafrost model into a large-scale ecosystem model: Evaluation of temporal and spatial scaling issues in simulating soil thermal dynamics. *J. Geophys. Res-Atmos.*, 106(D24), 33649-33670, 2001.
- 695 Zhuang, Q., McGuire, A. D., O'Neill, K. P., Harden, J. W., Romanovsky, V. E., & Yarie, J. Modeling soil thermal and carbon dynamics of a fire chronosequence in interior Alaska. *J. Geophys. Res-Atmos.*, 107(D1), FFR-3, 2002.



- 700 Zhuang, Q., McGuire, A. D., Melillo, J. M., Clein, J. S., Dargaville, R. J., Kicklighter, D. W., ... & Hobbie, J. E.
Carbon cycling in extratropical terrestrial ecosystems of the Northern Hemisphere during the 20th century: a
modeling analysis of the influences of soil thermal dynamics. *Tellus B.*, 55(3), 751-776, 2011.
- Zhuang, Q., Lu, Y., & Chen, M. An inventory of global N₂O emissions from the soils of natural terrestrial ecosystems.
Atmos. Environ., 47, 66-75, 2012.
- 705 Zhuang, Q., Chen, M., Xu, K., Tang, J., Saikawa, E., Lu, Y., ... & McGuire, A. D. (2013). Response of global soil
consumption of atmospheric methane to changes in atmospheric climate and nitrogen deposition. *Global
Biogeochem. Cy.*, 27(3), 650-663, 2013.



Table 1. Description of parameters used in the model

Parameters	Description	Unit	Reference Value	reference
N_fix	nitrogen fixation rate	g N m ⁻² day ⁻¹		
N_fixpot	potential nitrogen fixation rate	g N m ⁻² day ⁻²	0.01-1×10 ⁻³	Thornley (2001); Eckertsten et al.(2006); Corre-Hellou et al. (2007); Corre-Hellou et al. (2009);
ft	soil temperature factor	°C		
t_min	the minimum temperature for the start of N fixation	°C	0.5~5	Boote et al. (2008)
t_max	the maximum temperature for the stop of N fixation	°C	40~45	Boote et al. (2008)
t_optL	lower bond of optimal temperature	°C	10~20	Boote et al. (2008)
t_optH	upper bond of optimal temperature	°C	25~35	Boote et al. (2008)
fw	soil water factor			
φ1	coefficient for soil moisture		0	
φ2	coefficient for soil moisture		2	APSIM, EPIC (Sharpley and Williams, 1990; Boumiols et al., 1991; Cabelguenne et al., 1999); SOILN (Wu and McGechan, 1999)
Wa	lower bond of water content below which N fixation is totally restrict by the deficit of soil water		0	APSIM, EPIC (Sharpley and Williams, 1990; Boumiols et al., 1991; Cabelguenne et al., 1999); SOILN (Wu and McGechan, 1999)
Wb	upper bond of water content above which N fixation is not limited by the deficit of soil water		0.5	APSIM, EPIC (Sharpley and Williams, 1990; Boumiols et al., 1991; Cabelguenne et al., 1999)
Wf	available soil water content to that at field capacity			
f_Nup	parameter relating legume biological nitrogen fixation and soil nitrogen content		0.01~0.1	SOILN model (Wu and McGechan, 1999)
Ns	Soil mineral nitrogen content	g N m ²		
f_N	Soil mineral N effect			
f_C	Soil carbon effect			
Cr	Carbon concentration in the soil	g C g ⁻¹ soil		
Kc	Michaelis-Menten Constant for carbon	g C g ⁻¹ soil	0.001~0.01	Thornley (2001); Eckertsten et al. (2006)



Table 2. Calibration Sites of Biological Nitrogen Fixation Rate for Representative Ecosystems

Site name	Ecosystem Type	Lon.	Lat.	Experimental method	Reference	N Fixation Rate kg N ha ⁻¹ yr ⁻¹	Simulation, kg N ha ⁻¹ yr ⁻¹
Stordalen, Sweden	Tundra	18	68	ARA	Christie (1987); Sonesson et al. (1980)	2	2.5
Truelove Lowland, Canada	Tundra	-84.5	75.5	ARA	Chapin et al. (1990)	3	2.8
Niwot Ridge, Colorado, US	Tundra	-105.5	40	¹⁵ N	Bowman et al. (1996)	4.9	5.1
Central Sweden	Boreal Forest	18	60	ARA	Nohrstedt (1985)	0.93	0.9
PNFI, Ontario, Canada	Boreal Forest	-77	45.5	ARA	Hendrickson (1990)	0.25	1.2
Southern British Columbia, Canada	Boreal Forest	-119	49	ARA	Hendrickson and Burgess (1989)	2.8	2.2
Robson moraines, British Columbia	Boreal Forest	-119	53	ARA	Blundon and Dale (1990)	1.1	1.2
Umea, Sweden	Boreal Forest	19.5	64	ARA	Huss-Danell (1976)	1	1.5
Coweeta Basin,	Temperate Forest	-83	35	N accumulation	Boring and Swank (1984)	48	19.5
Hoh River, Washington, US	Temperate Forest	-123.5	48	ARA	Luken and Fonda (1983)	40	13
Tom Swamp, Massachusetts, US	Temperate Forest	-75	42.5	unspecified	Schwinzer (1983)	35	25.7
Big Creek Basin, Melbourne, Australia	Temperate Forest	145.5	38	ARA	Adams and Attwill (1984)	24	23.2
Jebo Creek, Utah, US	Temperate Forest	-112	42	¹⁵ N	Skujins et al. (1987)	10.2	12.5
Karri Forest, south-western Australia	Temperate Forest	116	-34.5	ARA	Grove and Malajczuk (1992)	7.93	8.5
Woodhill Forest, New Zealand	Temperate Forest	174.5	-37	N accumulation	Baker et al. (1986)	80	23.5
Gainesville, Florida, US	Temperate Forest	-82	30	N accumulation	Permar and Fisher (1983)	10.6	12.8
Fox park, Wyoming, US	Temperate Forest	-106	41	ARA	Fahey et al. (1985)	13	12.5
Mount Robson, Canada	Temperate Forest	-119	53.1	ARA	Blurdon and Dale (1990)	1.65	3.2
Dwellingup, South-western Australia	Temperate Forest	116	33	ARA	O'Connell and Grove (1987)	2.5	3.1
Adair, Oregon, US	Temperate Forest	-123	44.6	ARA	Heath et al (1988)	0.74	2.4



Priest River Experimental Forestry, Idaho, US	Temperate Forest	-116	48	ARA	Harvey et al (1989)	0.1	1.7
Arapaho Prairie, Nebraska, US	Grassland	-100	42	unspecified	Kaputsa and DuBois (1987)	0.2	0.7
Lynx Prairie Preserve, Ohio, US	Grassland	-83.5	39	ARA	DuBois and Kaputsa(1983)	8.2	1.9
Konza Prairie Research Natural Area, Kansas, US	Grassland	-96	39.5	nitrogenase activity	Eisele et al (1989)	21	3.3
Buso, Papua New Guinea	Tropical Forest	147	-7.5	ARA	Goosem and Lamb (1986)	0.5	5.2
Reserve Ducke, Manaus, Brazil	Tropical Forest	-59	-3	ARA	Sylvestre-Bradley et al. (1980)	2.45	3.5
Sinharaja Man and Biosphere reserve	Tropical Forest	80.5	6.5	ARA	Maheswaran and Gunatilleke (1990)	8	8.5
Amazon Territory of Venezuela	Tropical Forest	-67	2	ARA	Jordan et al (1983)	32	20.3
Kilauea, Hawaii, US	Tropical Forest	-155	19	ARA	Vitousek (1994)	2.8	18.5
Volcano La Soufriere, Guadeloupe	Tropical Forest	-61.5	16	ARA	Sheridan (1991)	4.02	7.2
Hawaii Volcanoes National Park, US	Tropical Forest	-155	19.5	ARA	Ley and D'Antonio (1998)	4.9	9.3
Santa Ynez Mountain, California, US	Mediterranean Shrubland	-120	34.5	ARA	Schlesinger et al.(1982)	1	2.4
San Bernardino Mountains, California, US	Mediterranean Shrubland	-116.5	34	ARA	Lepper and Fieschner (1977)	6.9	3.7
Harpers Well, California, US	Xeric Shrubland	-116	33.5	N accumulation	Rundel et al.(1982)	30	18.5
Sonoran Desert, Arizona, US	Xeric Shrubland	-112.5	33	cation accumulation	Jarrell and Virginia (1990)	40	23.5



Table 3. Sites used for model validation of N₂O emissions

Site name	Ecosystem		Lon.	Lat.	T (°C)	Precipitation (mm)	Length of experiment (day)	N ₂ O Fluxes kg N ha ⁻¹ yr ⁻¹	Reference
	Type	Temperate							
1 Germany	forest	Temperate	8.5	50.5	1~20	0-14	400	0.292	Müller and Shellock (2004)
2 Horquetas, Costa Rica	Savanna	Temperate	-85	10	25.8	3962	23-30	4.3	Veldkamp et al. (1998)
3 Yurimaguas, Peru	Rainforest	Temperate	-76	-6	26	2200	365	0.8	Palm et al. (2002)
4 Central Scotland	Grassland	Temperate	-4.5	56.5	12.6	129.3	7	0.02-0.09	Skiba et al. (1998)
5 Wagga Wagga	Grassland	Temperate	147.5	-35.5	3~31	570	90	0.35	Galbally et al. (2009)



Table 4. Model estimated biological nitrogen fixation in global natural terrestrial ecosystems

Ecosystem	Average coverage of N fixing plants	Coverage range	Reference	N Fixation Rate (kg N ha ⁻¹ yr ⁻¹)	Total Min (Tg N yr ⁻¹)	Total Max (Tg N yr ⁻¹)	Total Avg (Tg N yr ⁻¹)	Area (10 ⁸ ha)
wet tundra	9%	3%~15%	May and Webber (1982)	3.2	0.51	2.55	1.54	5.37
alpine tundra & wet tundra	9%	3%~15%	May and Webber (1982)	3.2	0.51	2.55	1.54	5.36
boreal forest	9%	4%~18%	Alexander and Billington (1986); weber and Van Cleve (1981)	2.1	2.01	9.06	4.53	19.3
temperate coniferous forest	5%	1%~10%	Cleveland et al (1999)	12.7	0.71	7.15	3.5	5.51
temperate deciduous forest	5%	1%~10%	Cleveland et al (1999)	12.7	0.76	7.65	3.75	5.89
temperate evergreen forest	5%	1%~10%	Cleveland et al (1999)	12.7	0.43	4.34	2.13	3.35
grassland	15%	5%~25%	Woodmansee et al (1981); Robertson and Rosswall (1986)	1.9	0.61	3.1	1.86	8.4
tropical forest	15%	5%~25%	Cleveland et al (2001)	18.2	10.8	54	32.6	17.8
xeric shrubland	15%	10%~20%	Johnson and Mayeux (1990)	5.7	2.92	14.6	8.35	14.8
Mediterranean shrubland	15%	10%~20%	Johnson and Mayeux (1990)	2.7	0.13	0.66	0.4	1.47
savanna	15%	5%~25%	Stewart et al (1978); Bate and Gunton (1982)	1.9	0.45	2.23	1.34	7.05
Total					19.84	107.89	61.54	94.3



Table 5. Model parameters for various natural terrestrial ecosystems

	N_{pot} (g N fixed day ⁻¹)	t_{optL} (°C)	t_{optH} (°C)	W_{upH} (J kg ⁻¹)	fNup	Kc (g C m ⁻²)
1 wet tundra	0.028	10	25	0.8	65	0.002
2 alpine tundra & wet tundra	0.028	10	25	0.8	65	0.002
3 boreal forest	0.032	12	25	0.8	70	0.008
4 temperate coniferous forest	0.55	16	35	0.6	80	0.01
5 temperate deciduous forest	0.55	18	35	0.6	80	0.01
6 temperate evergreen forest	0.55	18	35	0.6	80	0.01
7 grassland	0.05	18	35	0.5	60	0.012
8 tropical forest	0.8	20	35	0.8	100	0.005
9 xeric shrubland	0.7	15	35	0.4	65	0.016
10 Mediterranean shrubland	0.08	19	35	0.5	65	0.016
11 savanna	0.05	20	35	0.5	60	0.012





Figure 1. Sites used with observation data of N₂ fixation rate (Blue) and N₂O emissions (Red) for model parameterization



(a)



(b)






Figure 2. Model sensitivity of N fixation and N₂O emissions in natural terrestrial ecosystems to changing model input data: Increasing or decreasing each variable by 10% for air temperature (T), precipitation (P), soil carbon content (C), soil nitrogen content (N) for (a) N fixation rate and (b) N₂O emissions



(a)



(b)



(c)





Figure 3. Comparison between Modeled and Observed Nitrogen Fixation Rate at site level: (a) All sites, (b) Temperate Forest, (c) Tropical Forest (data listed in Table 2). Y is simulated N fixation while X represents the observational N fixation.



(a)



(b)






Figure 4. Simulated spatial distribution of (a) BNF rates ($\text{kg N ha}^{-1} \text{ yr}^{-1}$); (b) N_2O emissions ($\text{kg N ha}^{-1} \text{ yr}^{-1}$) in natural terrestrial ecosystems during 1990-2000 by considering the BNF effects.



(a)




(b)






(c)



(d)





(e)




Figure 5. Comparison between simulated and observed N_2O emissions ($\text{g N m}^{-2} \text{ day}^{-1}$) with two versions of the model (Red indicates the simulations considering BNF and blue indicates the simulations without considering BNF effects on N_2O emissions) at sites: (a) Temperate Forest (50.5°N , 8.5°E); (b) Tropical Forest (10°S , 63°W); (c) Savanna (10°N , 85°W); (d) Tropical Forest (6°S , 76°W).